

REPLACING CORNMEAL WITH MANGO MEAL IN DIETS FOR JUVENILE TAMBAQUI *Colossoma macropomum*: GROWTH AND METABOLIC PARAMETERS

Anderson Miranda de SOUZA¹

Daniela Ferraz Bacconi CAMPECHE²

Gilberto MORAES³

Fúlvio Viegas Santos Teixeira de MELO⁴

Manoel Adriano da Cruz NETO⁵

José Fernando Bibiano MELO⁵

ABSTRACT

The effect on tambaqui fingerlings performance caused by diets containing two different carbohydrate sources was evaluated. Four isoproteic diets with different mango meal concentrations (0, 33, 66 and 100%) replacing cornmeal were tested. The feeding trial lasted 45 days and the feeding rate was 8% of total biomass. 240 tambaqui fingerlings with an initial weight average of 3.66g were distributed in 12 500-liter tanks in a recirculating water system with biofilter. Performance, feed intake, carcass yield, survival, total cholesterol, plasmatic free amino acids, hepatic aspartate aminotransferase (AST), hepatic glycogen and plasma glucose values were evaluated. The results showed that weight gain, daily weight gain, specific growth rate, apparent food consumption, cholesterol, AST and glycogen showed higher values in the treatments with the highest levels of mango meal. Apparent feed conversion, carcass yield, survival and plasma glucose were not affected by the two sources of carbohydrates tested. In conclusion, mango meal used in a practical diet for tambaqui improved growth performance.

Key words: carbohydrate; performance; fish nutrition; metabolic profile.

¹Universidade Federal da Bahia – UFBA, Rua Barão de Jeremoabo, 668, Ondina, CEP 40170-115, Salvador, BA, Brasil.

²Embrapa Semiárido, Rodovia BR-428, Km 152, s/n, Zona Rural, CEP 56302-970, Petrolina, PE, Brasil.

³Universidade Federal de São Carlos – UFSCar, Rodovia Washington Luís, Km 235, SP-310, CEP 13565-905, São Carlos, SP, Brasil.

⁴Instituto Federal Baiano – IFBA, Campus Catu, Rua Matias Inácio Peneluc, 118, Centro, CEP 48110-000, Catu, BA, Brasil.

⁵Universidade Federal do Vale do São Francisco – UNIVASF, Rodovia BR-407, Km 12, Lote 543, s/n, Projeto de Irrigação Nilo Coelho, CEP 56300-000, Petrolina, PE, Brasil. E-mail: melojfb@yahoo.com.br (corresponding author).

SUBSTITUIÇÃO DO FARELO DE MILHO POR FARINHA DE MANGA EM DIETAS PARA TAMBAQUI *Colossoma macropomum* JUVENIL: CRESCIMENTO E PARÂMETROS METABÓLICOS

RESUMO

Avaliou-se o efeito sobre o desempenho de alevinos de tambaqui alimentados com dietas contendo duas fontes diferentes de carboidratos. Foram testadas quatro dietas isoprotéicas com diferentes concentrações de farinha de manga (0, 33, 66 e 100%) em substituição ao farelo de milho. O teste de alimentação foi de 45 dias e a taxa de alimentação foi de 8% da biomassa total. Foram utilizados 240 alevinos de tambaqui com peso inicial médio de 3,66g distribuídos em 12 caixas de 500L em sistema de recirculação de água com biofiltro. Foram avaliados os índices de desempenho, consumo de ração, rendimento de carcaça, sobrevivência, colesterol total, aminoácidos livres plasmáticos, aspartato aminotransferase hepática (AST), glicogênio hepático e glicose plasmática. Os resultados mostraram que o ganho ponderal de peso, a taxa de crescimento específico, o consumo aparente de dieta, colesterol, AST e glicogênio apresentaram valores maiores nos tratamentos com os maiores níveis de farinha de manga. A conversão alimentar aparente, o rendimento de carcaça, a sobrevivência e a glicose plasmática não foram afetados pelas duas fontes de hidratos de carbono testadas. Em conclusão, a farinha de manga utilizado na dieta prática para tambaqui melhorou o desempenho.

Palavras-chave: carboidrato; desempenho; nutrição de peixes; perfil metabólico.

Received: May 24, 2017

Approved: March 06, 2018

INTRODUCTION

The continuous demand for fish, not only due to demographic growth but also as a healthy food preference has made fish production viable, as have the development of new technologies related to it.

Nutrition has been a technology tool that helps to improve fish growth in a shorter time and contributes to the benefit of worldwide production. Numerous ingredients have been tested around the world in different fish species (XAVIER *et al.*, 2012; PEREIRA JUNIOR *et al.*, 2013; MUIN *et al.*, 2013; NAQVI *et al.*, 2015). There is a

great variety of vegetable-source ingredients with the potential to be used in fish feed (GABER, 2006; SAMPAIO-OLIVEIRA and CYRINO, 2008; NYINA-WAMWIZA *et al.*, 2010; PEREIRA JUNIOR *et al.*, 2013), since many species consume vegetables as their natural feeding behavior (LUCAS, 2008). According to BATISTA *et al.* (2015), mango (*Mangifera indica* L.) is rich in carbohydrates such as starch, reducing sugars and pectic substances. This fruit is very important for Brazil and for exports, however there is a large surplus volume generating food waste, environmental contamination and economic loss. According to MARQUES *et al.* (2010) the mango has in its composition lipids (0.61% ± 0.03), proteins (0.44% ± 0.08), non-reducing sugars (4.13% ± 0.12), starch (0.15% ± 0.16), dietary fiber (3.28% ± 0.28), and total carbohydrates (16.5%). Studies have been done testing the fruit or its by-products for its use in fish feed and its effects on performance and metabolism (SILVA *et al.*, 2003; GUIMARÃES and STORTI-FILHO, 2004; AZAZA *et al.*, 2009; SOUZA *et al.*, 2013). Therefore, it can be an important source of feed ingredients in the future and is already proving to be a successful ingredient (BEZERRA *et al.*, 2014) since it has a carbohydrate profile that is used by omnivore fish (KANSCI *et al.*, 2003).

Intermediary metabolism parameters in the plasma and liver have been used as important tools to evaluate the effects of carbohydrates in fish diets (SKIBA-CASSY *et al.*, 2013). Plasma glucose is an indicator of the ingested saccharide effect on fish metabolism (BOOTH *et al.*, 2013). Total free amino acids and growth performance parameters together may indicate if dietary protein has been mobilized as a key mechanism in metabolism adaptation (COSTAS *et al.*, 2011). It has also been used to evaluate the effect of alternative ingredients for fish feed (ADAMIDOU *et al.*, 2009; HANSEN *et al.*, 2011; AZEVEDO *et al.*, 2013).

Brazil produced 1.163 million tons of mangoes in 2013, most notably in the states of Bahia, São Paulo and Pernambuco, which together accounted for 77.25% of Brazilian production. With these results, Brazil is the seventh largest producer and the fourth largest exporter of mango in the world (AZERÉDO *et al.*, 2016). This situation might be an opportunity for its use as an ingredient in animal feed, since mango and its by-products are a source of fiber and carbohydrates (RÉGO *et al.*, 2010; PEREIRA JUNIOR *et al.*, 2013). In *in natura* and dried mangos, total sugar can reach values of 52.49%, where 17% can be glucose, and fructose values can vary from 2.3 to 3.1% (BERNARDES-SILVA *et al.*, 2003). It is known that fructose reaches the cell first, since it does not need to pass through an enzymatic pathway (ENES *et al.*, 2009). This advantage can possibly make fish grow faster, with less energy costs.

This work aims to evaluate juvenile tambaqui growth performance and metabolic profile, when fed different inclusion levels of mango meal as a replacement for cornmeal.

METHODS

The trial was carried out in the Aquiculture Laboratory from the Federal University of Vale do São Francisco in Petrolina-PE, Brazil, through the ethics committee protocol 0016/14415 on animal use.

Experimental diets

Four experimental diets were formulated, replacing cornmeal with mango meal: 0; 33; 66 and 100%. Diets were prepared according to the species' nutritional requirements (RODRIGUES, 2016) at the age studied, while performing four experimental diets (Table 1).

Mango meal was made in the university laboratory with mango pulp, cut in small pieces and taken to a forced air circulation oven at 65 °C for 72 hours. When it came out it was crushed in a mill and stored frozen at -18 °C until the feeds were made. The other feed ingredients were also crushed in a mill and sifted on a 0.5 mm net. All ingredients were hand mixed to complete homogenization. Water at 50 °C was added to the mixture. Pellets were produced in a pellet mill and then dried in a forced air circulation oven at 65 °C for 24 hours. Pellets were ground in a manual mill and separated in different diameters with different-sized net until it had the specific diameter for fish size in accord with their development. Diet composition was analyzed as established by the Association of Official Analytical Chemists (AOAC, 2000).

Table 1. Formulation and chemical composition of the experimental diets.

Ingredients (g kg ⁻¹)	Experimental diets			
	0%	33%	66%	100%
Soybean meal	621.2	621.2	621.2	621.2
Cornmeal	300.0	200.0	100.0	0
Mango meal ¹	0	99.9	199.9	300.0
Bicalcic phosphate	30.3	30.3	30.3	30.3
Soybean oil	22.3	22.3	22.3	22.3
Premix ²	20.0	20.0	20.0	20.0
NaCl	5.0	5.0	5.0	5.0
DL-methionine	1.0	1.0	1.0	1.0
Butylhydroxy-toluene	00.1	00.1	00.1	00.1
<i>Chemical composition</i>				
<i>(g kg⁻¹)*</i>				
Dry matter	940.0	952.4	936.6	933.8
Crude protein	348.0	336.3	334.1	329.9
Lipid	34.6	33.4	32.2	31.0
Total Carbohydrates ³	270.7	265.0	259.3	253.6
Mineral matter	78.3	79.8	81.0	82.2
Crude energy (Kcal kg ⁻¹)	3.261	3.290	3.319	3.339

¹Composition in g kg⁻¹: 93.51 dry matter (%); 2.81 ash (%); 1.35 lipid (%); 3.99 crude protein (%) and gross energy is 4619.04 Kcal kg⁻¹. ²A=1,200,000 UI; D3=200,000 UI; E = 12,000 mg; K3 = 2400 mg; B1 = 4800 mg; B2 = 4800 mg; B6 = 4000 mg; B12 = 4800 mg; folic acid = 1200 mg; calcium pantothenate = 12,000 mg; vit. C = 48,000 mg; biotin = 48 mg; choline = 65,000 mg; nicotinic acid = 24,000 mg; minerals: Fe = 10,000 g; Cu = 600 mg; Mn = 4000 mg; Zn = 6000 mg; I = 20 mg; Co = 2 mg e Se = 20 mg. ³*Calculated value based on dry matter (SUPERCRAC 2006).

Experimental design

Fish were stocked in 12 500-liter plastic tanks. Tanks were organized in a closed recirculation system with a biofilter, with physical and biological treatments. Twenty juvenile tambaqui with an initial weight of 3.66 ± 0.84 g and initial length of 5.79 ± 0.50 cm were placed in each tank and acclimated for 15 days before the trial began and were fed the controlled diet.

Organic matter was daily removed from the tank's bottom by suction. Water parameters of dissolved oxygen, temperature and pH were daily monitored with automatic monitors. Water quality parameters were analyzed daily with a multiparameter probe. The average values were: 27.12 ± 1.31 temperature; 5.4 ± 1.08 mg L⁻¹ for dissolved oxygen; 0.13 ± 0.001 mS for electrical conductivity; 7.77 ± 1.24 for pH; and 0.01 ± 0.001 mg L⁻¹ ammonia (NH₃). During the experiment these parameters were within the recommendations for the culture of this species (SIPAÚBA-TAVARES, 1995).

Feed was offered at 8% of fish biomass, twice a day, at 9:00 a.m. and 5:00 p.m., with the experimental diets performing four treatments in triplicate. Every 10 days fish were weighed to adjust these values, but the last weighing was done within 5 days. The experiment was performed over a 45-day period.

Growth parameters in each treatment were calculated by analyzing: weight gain (WG, g) = final body weight – initial body weight; length gain (LG, cm) = final body length – initial body length; daily weight gain (DWG, g) = f(final body weight – initial body weight)/days of trial; specific growth rate (SGR %) = $100 \times (\ln \text{ final weight} - \ln \text{ initial weight})/\text{days of trial}$; carcass yield (CY, g) = final body weight – viscera weight; feed intake (FI, g) = feed offered related to body weight % x days of trial; apparent feed conversion (AFC) = feed intake/weight gain; and survival (%) [S = $100 \times (\text{initial number of fish} - \text{final number of fish})$]. For performance evaluation, 240 fish dispersed in the experimental units were measured at the beginning and at the end of the trial.

Metabolic analyses

At the end of the experimental period, 10 animals from each treatment were sampled for blood collection and were considered as repetition. Fish were anesthetized with benzocaine (1g 10L⁻¹)

and blood was withdrawn through the caudal vein, with heparinized syringes. Afterwards, fish were euthanized by medullar section for liver collection. Metabolite concentration analyses were carried out in plasma and tissue extracts.

Free amino acids (mg dL⁻¹) were determined in neutral extracts at 570 nm (COPLEY, 1941). Liver glycogen ($\mu\text{mol glucose mg/ hepatic protein}$) was assayed in alcoholic precipitates from alkaline tissue homogenates (BIDINOTTO *et al.*, 1997; DUBOIS *et al.*, 1960).

Plasma glucose was quantified by an enzymatic method using a commercial kit with the reading done at a 340 nm wavelength. 10 microliters of sample incubated for 5 minutes at 37 °C were used. Total plasmatic cholesterol (mg dL⁻¹) was determined by a colorimetric enzymatic method, using a commercial kit with the reading done at a 500 nm wavelength. Ten (10) microliters of plasma sample was incubated for 10 minutes at 37 °C.

Aspartate Amino Transferase (AST) enzymatic activity (U mg⁻¹ protein) in the homogenized liver samples was determined by a kinetic method where the absorbance reduction in 340 nm was registered every 2 minutes. 30 microliters of homogenized liver were used. Enzymes were determined by a kinetic method (KANEKO, 1989) using a commercial kit. Two readings were performed in two minutes taking the average of three readings with a 60 second gap, using a 340 nm filter. The total protein was analyzed in plasma and liver with a commercial kit. 10 microliters of plasma and 100 mg of homogenized liver in pH 7.0 buffer were used. The samples were read at 545 nm.

Statistics

The experimental design was established as incomplete blocks formed by four treatments and ten repetitions per treatment. Metabolite and enzyme data were submitted to variance analysis. The Tukey test ($p < 0.05$) was applied to measure average difference in the parameters among different treatments.

RESULTS

Growth parameters

Weight gain (WG) and daily weight gain (DWG) showed significant differences among treatments (Table 2). It can be observed that juveniles fed a diet containing 100% of mango meal

Table 2. Growth performance of *Colossoma macropomum* juvenile fed diet containing increasing levels of mango meal replacing cornmeal.

Performance	Diets			
	0%	33%	66.0%	100%
Weight Gain (g fish ⁻¹)	11.97 ^b ±1.60	12.91 ^b ±1.77	15.12 ^{ab} ±2.08	17.59 ^a ±1.98
Daily weight gain (g fish ⁻¹)	0.26 ^b ±0.01	0.29 ^b ± 0.01	0.33 ^{ab} ±0.01	0.39 ^a ±0.009
Specific growth rate (%)	26.59 ^b ±3.21	28.70 ^b ±2.54	33.59 ^{ab} ±4.12	39.10 ^a ±3.99
Feed consumption (g fish ⁻¹)	16.14 ^b ±2.10	17.24 ^{ab} ±3.22	20.05 ^a ±2.30	20.31 ^a ±2.16
Feed conversion rate	1.36	1.34	1.33	1.16
Carcass yield (%)	80.78±3.77	85.22±3.90	83.89±4.18	79.98±3.61
Survival (%)	100	100	100	100

Metabolic profile.

Table 3. Metabolic intermediates and enzymatic activity in *Collossoma macropomum* juvenile fed diets containing increasing levels of mango meal replacing cornmeal.

	Tissue Variable	Diets			
		0%	33%	66%	100%
Plasma	Cholesterol (mg dL ⁻¹)	104.84 ^a ±12.33	44.09 ^b ±4.21	66.13 ^b ± 5.62	52.42 ^b ±5.00
	Free amino acids (µmols dL ⁻¹)	10.61 ^c ±0.98	37.55 ^a ±2.12	16.03 ^{bc} ±1.55	20.72 ^b ±2.34
	Glucose (mg dL ⁻¹)	68.5±3.70	75.8±4.09	71.0±3.86	63.8±3.43
Liver	AST (U mg ⁻¹ protein)	6.39 ^b ±0.95	10.65 ^{ab} ±0.90	12.60 ^{ab} ±1.03	16.36 ^a ±0.88
	Glycogen (µmol glucose mg ⁻¹ g tissue)	37.28 ^a ±2.76	19.19 ^b ±2.90	23.26 ^{ab} ±3.16	22.00 ^{ab} ±3.69

showed higher growth, with an average value of 17.59 g for WG and 0.39 g.day⁻¹ for DWG. The specific growth rate also showed a significant difference among evaluated treatments ($p < 0.05$) following the same patterns of the other evaluated parameters for fish fed a diet with 100% of mango meal. Feed consumption was also influenced by the mango meal percentage in the diet. The highest values were registered among treatments with higher inclusion of mango meal.

Feed conversion rate, carcass composition and survival were not influenced by levels of mango meal inclusion in the diets. There was no significant difference among treatments related to these parameters.

Increasing replacement of cornmeal by mango meal significantly reduced plasmatic cholesterol, increased free amino acids and increased AST activity in the liver (Table 3). Plasma glucose was kept constant in all treatments. There was a significant hepatic glycogen deposit in the treatment's control group.

DISCUSSION

Weight gain in juvenile fish is indicative of feed's nutritional adequacy (CHO *et al.*, 1986). In this trial, the diet containing a higher proportion of mango meal provided higher weight gain in the fish. This fact must be due to the high amounts fructose present in the mango meal as a simple carbohydrate source (KANSCI *et al.*, 2003). Simple carbohydrate sources, such as fructose, demand less metabolic energy to provide energy to the animal body. Fruit and fruit by-products have been tested in fish feed with good performance results at different replacement levels (FALAYE and OLORUNTUYI, 1998; ULLOA ROJAS and VERRETH, 2003; LOCHMANN *et al.*, 2009; CAMPECHE *et al.*, 2014).

Cornmeal replacement by millet and mesquite for tambaqui juvenile diets, did not show significant differences among treatments for weight gain and feed consumption (MIRANDA *et al.*, 2009; SILVA *et al.*, 1997), which is contrary to this study. However, like millet and mesquite, mango meal in tambaqui diet did not affect feed conversion rate. Juvenile tambaqui growth performance was not affected when cornmeal was replaced by pupunha meal (*Bactris gasipaes*) in the diet (MORI-PINEDO *et al.*, 1999). The same was observed when pupunha (*Bactris gasipaes*) replaced wheat bran in feeds for tambaqui (LOCHMANN *et al.*, 2009). This fact must be due to the carbohydrate source found in the

different diets evaluated as well due to the tambaqui feeding plasticity (DABROWSKI and PORTELLA, 2006). The effects of alternative carbohydrate sources on growth parameters might also depend on its inclusion percentage (NAGAE *et al.*, 2002; SIGNOR *et al.*, 2007). A significant variety of items can be part of an Amazon fish's natural diet, among them fruits and seeds that are natural sources of all kinds of nutrients (SILVA *et al.*, 2003). According to LOVELL (1989) high water temperature promotes growth rate improvement as it maintains metabolic activities constant allowing energy ingested from the feed to be used for growth. Therefore, tropical species can show better feed conversion rates. Although tilapia and tambaqui have different food habits and may belong to different habitat climates, both accept very well a great diversity of feed ingredients.

Carcass yield can help to indicate how nutrients ingested were utilized by the animal. In the present study, although a difference in growth was observed, carcass yield did not show statistical differences, which is similar to what was observed for tilapia fed with cassava meal replacing cornmeal (BOSCOLO *et al.*, 2002). In these cases, feed protein and energy efficiency analyses associated to metabolic parameters help to understand feed nutrients used by fish. Complex carbohydrates can be well used by fish when pre-treated to make its digestibility easier and better (DAVIES *et al.*, 2011; RENUKDAS *et al.*, 2014).

Plasmatic cholesterol reduction obtained in this study may be due to the carbohydrate source chosen, which was mango meal. There are two hypotheses related to the reduction effect mechanism in plasmatic cholesterol. The first is that soluble fiber found in fruits, vegetables and oats, for instance, increase bile acids leading the liver to remove cholesterol from the blood (DE ANGELIS, 2001). The other hypothesis shows that propionate, which is the product from soluble fiber fermentation, inhibits hepatic cholesterol synthesis. Although there are some controversies in the mechanism related to fibers, the plasmatic cholesterol reduction hypothesis is becoming strong (GALLAHER and SCHNEEMAN, 1997).

Glycaemia maintenance was efficient even with the change in the carbohydrate source. Glycaemia levels can indicate the utilization of carbohydrate source and inclusion percent by a fish species (BOOTH *et al.*, 2013). Although a reduction in the hepatic glycogen seems to be related to glycaemia in order to keep its values; as was found in this study. The difference in glycogen levels was also observed in *Collossoma macropomum* and *Piaractus brachypomus* fed carbohydrate-rich alternative

foodstuffs with 30% inclusion (LOCHMANN *et al.*, 2009). The metabolic profile shown in the plasma and liver is an adaptation to keep energy maintenance. In fact, nutrient sources and concentrations can demonstrate different responses from what was registered in this study. European seabass (*Dicentrarchus labrax*), a carnivorous species fed diets with different legumes as an alternative carbohydrate source, did not show any difference in hepatic glycogen index (ADAMIDOU *et al.*, 2009). This difference from results found in the present work may be related to the natural feeding habitat difference.

Since the increase in plasma free amino acids can be related to growth improvement or lack of nutrients (COSTAS *et al.*, 2011), it is a very important variable to be observed. The difference in feed protein or energetic ingredients clearly affects plasma free amino acids (MACH and NORTVEDT, 2011; LARSEN *et al.*, 2012). This fact is a consequence of the ingredient's amino acid profile and affects fish growth. Plasma free amino acid also leads to the better use of the carbohydrate source (mango meal) for the energetic process and, as a consequence, the use of available amino acids for fish growth. This, due to a possible sparing effect of dietary protein, is for maintaining observed glucose and increased amino acid levels in plasma. In this study, better growth was observed in the fish group with a higher inclusion of mango meal. Since the feed was isonitrogenous it was clearly noticed that plasma free amino acid differences were due to its different metabolic use as a consequence of the carbohydrate sources used in the feed. Available amino acids might be deaminated and excreted, or used as energy which might explain the AST activity increase. MOYANO *et al.* (1991) suggests that transaminases and deaminase quantification might be a metabolic indication of amino acid excess in fish. The increase of protein and the reduction of carbohydrate concentrations in *Leiarius marmoratus*'s diet led to an increase in transaminase activity, suggesting gluconeogenesis (SOUZA *et al.*, 2014). Surplus amino acids required for protein biosynthesis are deaminated and carbon residues are converted in the Krebs cycle into fats or carbohydrates (DRIEDZIC and HOCHACHKA, 1978). This metabolic pathway of amino acid biosynthesis for energy production was verified in Jundia (*Rhamdia quelen*) fed high protein concentrations (MELO *et al.*, 2016). The use of fruit for fish feed is still a recent practice. Further studies should be performed to verify the effects of mango from the fractions that compose this fruit, together with studies of digestibility to prove its efficiency.

In conclusion, data from this trial show that tambaqui growth parameters such as weight gain, and specific growth rate were improved. In addition, the replacement did not cause physiological injuries or affect feed conversion rates and carcass yield. Mango meal possesses an optimal metabolic profile for energy use that comes from its carbohydrate source as was tested in tambaqui.

REFERENCES

ADAMIDOU, S.; NENGAS, I.; HENRY, M.; GRIGORAKIS, K.; RIGOS, G.; NIKOLOPOULOU, D.; KOTZAMANIS, Y.; BELL, G.J.; JAUNCEY, K. 2009 Growth feed utilization: health and organoleptic characteristics

- of European seabass (*Dicentrarchus labrax*) fed extruded diets including low and high levels of three different legumes. *Aquaculture*, 293(3-4): 263-271. <http://dx.doi.org/10.1016/j.aquaculture.2009.04.045>.
- ANGELIS, R.C. 2001 *Conceitos de nutrientes não tradicionais: importância de alimentos vegetais na proteção da saúde*. Belo Horizonte: Atheneu. 75p.
- AOAC – Association of Official Analytical Chemistry. 2000 *Official methods of analysis*. 17th ed. Washington: AOAC.
- AZAZA, M.S.; MENSİ, F.; KAMMOUN, W.; ABDELOUAHEB, A.; BRINI, B.; KRAÏEM, M. 2009 Nutritional evaluation of waste date fruit as partial substitute for soybean meal in practical diets of juvenile Nile tilapia *Oreochromis niloticus* L. *Aquaculture Nutrition*, 15(3): 262-272. <http://dx.doi.org/10.1111/j.1365-2095.2008.00591.x>.
- AZERÊDO, L.P.M.; SILVA, S.M.; LIMA, M.A.C.; DANTAS, R.L.; PEREIRA, W.E. 2016 Qualidade de manga 'tommy atkins' da produção integrada recoberta com fécula de mandioca associada a óleos essenciais e quitosana. *Revista Brasileira de Fruticultura*, 38(1): 141-150. <http://dx.doi.org/10.1590/0100-2945-270/14>.
- AZEVEDO, R.V.; TONINI, W.C.T.; BRAGA, L.G.T. 2013 Óleo e torta de dendê em rações para juvenis de tilápia-do-nilo. *Pesquisa Agropecuária Brasileira*, 48(8): 1028-1034. <http://dx.doi.org/10.1590/S0100-204X2013000800031>.
- BATISTA, P.F.; LIMA, M.A.C.D.; TRINDADE, D.C.G.D.; ALVES, R.E. 2015 Quality of different tropical fruit cultivars produced in the Lower Basin of the São Francisco Valley. *Revista Ciência Agrônômica*, 46(1): 176-184. <http://dx.doi.org/10.1590/S1806-66902015000100021>.
- BERNARDES-SILVA, A.P.F.; LAJOLO, F.M.; CORDENUNSI, D.R. 2003 Evolução dos teores de amido e açúcares solúveis durante o desenvolvimento e amadurecimento de diferentes cultivares de manga. *Food Science and Technology*, 23(3): 116-120.
- BEZERRA, K.S.; SOUZA, R.C.; MELO, J.F.B.; CAMPECHE, D.F.B. 2014 Crescimento de tambaqui alimentado com diferentes níveis de farinha de manga e proteína na ração. *Archivos de Zootecnia*, 63(244): 587-598. <http://dx.doi.org/10.4321/S0004-05922014000400003>.
- BIDINOTTO, P.M.; SOUZA, R.H.S.; MORAES, G. 1997 Hepatic glycogen in eight tropical freshwater teleost fish: a procedure for field determinations of microsamples. *Boletim Técnico CEPTA*, 10: 53-60.
- BOOTH, M.A.; MOSES, M.D.; ALLAN, G.L. 2013 Utilization of carbohydrate by yellowtail kingfish *Seriola lalandi*. *Aquaculture*, 376: 151-161. <http://dx.doi.org/10.1016/j.aquaculture.2012.11.024>.
- BOSCOLO, W.R.; HAYASHI, C.; MEURER, F. 2002 Farinha de Varredura de Mandioca (*Manihot esculenta*) na Alimentação de Alevinos de Tilápia do Nilo (*Oreochromis niloticus* L.). *Revista Brasileira de Zootecnia*, 31(2): 546-551. <http://dx.doi.org/10.1590/S1516-35982002000300002>.
- CAMPECHE, D.F.B.; MELO, J.F.B.; BALZANA, L.; SOUZA, R.C.; FIGUEIREDO, R. 2014 Farelo de licuri em dietas para alevinos de tambaqui (*Colossoma macropomum*. Cuvier. 1818). *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 66(2): 539-545. <http://dx.doi.org/10.1590/1678-41625920>.
- CHO, C.Y.; COWEY, C.B.; WATANABE, T. 1986 Finfish nutrition in Asia: methodological approaches to research and development. *International Development Research Center*, 233: 154.
- COPLEY, N.G. 1941 Alloxan and ninhydrin test. *Analyst*, 66: 492-493.

- COSTAS, B.; ARAGÃO, C.; RUIZ-JARABO, I.; VARGAS-CHACOFF, L.; ARJONA, F.J.; DINIS, M.T.; MANCERA, J.M.; CONCEIÇÃO, L.E.C. 2011 Feed deprivation in Senegalese sole (*Solea senegalensis* Kaup. 1858) juveniles: effects on blood plasma metabolites and free amino acids levels. *Fish Physiology and Biochemistry*, 37(3): 495-504. <http://dx.doi.org/10.1007/s10695-010-9451-2>. PMID:21107905.
- DABROWSKI, K.; PORTELLA, M.C. 2006 Feeding plasticity and nutritional physiology in tropical fishes. In: VAL, A.L.; ALMEIDA-VAL, V.M.F.D.; RANDALL, D.J. *The physiology of tropical fishes*. Amsterdam: Elsevier. p. 155-224. (Fish Physiology, 21).
- DAVIES, S.J.; ABDEL-WARITH, A.A.; GOUVEIA, A. 2011 Digestibility characteristics of selected feed ingredients for developing bespoke diets for Nile tilapia culture in Europe and North America. *Journal of the World Aquaculture Society*, 42(3): 388-398. <http://dx.doi.org/10.1111/j.1749-7345.2011.00478.x>.
- DRIEDZIC, W.R.; HOCHACHKA, P.W. 1978 Metabolism in fish during exercise. In: HOAR, W.S.; RANDALL, D.J. *Fish physiology*. New York: Academic Press. v. 7. 543p.
- DUBOIS, M.G.; GILLES, K.A.; HAMILTON, J.K.; REBERS, P.A.; SMITH, F. 1960 Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, 28(3): 350-358. <http://dx.doi.org/10.1021/ac60111a017>.
- ENES, P.; PANSERAT, S.; KAUSHIK, S.; OLIVA-TELES, A. 2009 Nutritional regulation of hepatic glucose metabolism in fish. *Fish Physiology and Biochemistry*, 35(3): 519-539. <http://dx.doi.org/10.1007/s10695-008-9259-5>. PMID:18791853.
- FALAYE, A.E.; OLORUNTUYI, O.O. 1998 Nutritive potential of plantain peel meal and replacement value for maize in diets of African catfish (*Clarias gariepinus*) fingerlings. *Tropical Agriculturist*, 75(4): 488-492.
- GABER, M.M. 2006 The effects of plant-protein-based diets supplemented with yuca on growth digestibility and chemical composition of Nile tilapia (*Oreochromis niloticus*. L.) fingerlings. *Journal of the World Aquaculture Society*, 37(1): 74-81. <http://dx.doi.org/10.1111/j.1749-7345.2006.00008.x>.
- GALLAHER, D.D.; SCHNEEMAN, B.O. 1997 Fibra alimentaria. In: BOWMAN, B.; RUSSELL, R. *Conocimientos actuales sobre nutrición*. Washington: International Life Sciences Institute Press. p. 95-105.
- GUIMARÃES, S.F.; STORTI-FILHO, A. 2004 Produtos agrícolas e florestais como alimento suplementar de tambaqui em policultivo com jaraqui. *Pesquisa Agropecuária Brasileira*, 39(3): 293-296. <http://dx.doi.org/10.1590/S0100-204X2004000300014>.
- HANSEN, J.O.; SHEARER, K.D.; OVERLAND, M.; PENN, M.H.; KROGDAHL, Å.; MYDLAND, L.T.; STOREBAKKEN, T. 2011 Replacement of LT fish meal with a mixture of partially deshelled krill meal and pea protein concentrates in diets for Atlantic salmon (*Salmo salar*). *Aquaculture*, 315(3): 275-282. <http://dx.doi.org/10.1016/j.aquaculture.2011.02.038>.
- KANEKO, J.J. 1989 *Clinical biochemistry of domestic animals*. 4th ed. San Diego: Academic. 932p.
- KANSCI, G.; KOUBALA, B.B.; LAPE, I.M. 2003 Effect of ripening on the composition and the suitability for jam processing of different varieties of mango (*Mangifera indica*). *African Journal of Biotechnology*, 2(9): 296-300.
- LARSEN, B.K.; DALSGAARD, J.; PEDERSEN, P.B. 2012 Effects of plant proteins on postprandial, free plasma amino acid concentrations in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 326: 90-98. <http://dx.doi.org/10.1016/j.aquaculture.2011.11.028>.
- LOCHMANN, R.; CHEN, R.; CHU-KOO, F.W.; CAMARGO, W.N.; KOHLER, C.C.; KASPER, C. 2009 Effects of carbohydrate-rich alternative feedstuffs on growth, survival, body composition, hematology and non-specific immune response of black pacu. *Colossoma macropomum* and red pacu. *Piaractus brachypomus*. *Journal of the World Aquaculture Society*, 40(1): 33-44. <http://dx.doi.org/10.1111/j.1749-7345.2008.00232.x>.
- LOVELL, T. 1989 *Nutrition and feeding of fish*. New York: Van Nostrand Reinhold. 260p. <http://dx.doi.org/10.1007/978-1-4757-1174-5>.
- LUCAS, C.M. 2008 Within flood season variation in fruit consumption and seed dispersal by two characin fishes of the Amazon. *Biotropica*, 40(5): 581-589. <http://dx.doi.org/10.1111/j.1744-7429.2008.00415.x>.
- MACH, D.T.N.; NORTVEDT, R. 2011 Free amino acid distribution in plasma and liver of juvenile cobia (*Rachycentron canadum*) fed increased levels of lizardfish silage. *Aquaculture Nutrition*, 17(2): 644-656. <http://dx.doi.org/10.1111/j.1365-2095.2010.00814.x>.
- MARQUES, A.; CHICAYBAM, G.; ARAUJO, M.T.; MANHÃES, L.R.T.; SABAA-SRUR, A.U. 2010 Composição centesimal e de minerais de casca e polpa de manga (*Mangifera indica* L.) cv. Tommy atkins. *Revista Brasileira de Fruticultura*, 32(4): 1206-1210. <http://dx.doi.org/10.1590/S0100-29452010005000117>.
- MELO, J.F.B.; LUNDSTEDT, L.M.; INOUE, L.A.K.; METÓN, I.; BAANANTE, I.; MORAES, G. 2016 Glycolysis and gluconeogenesis in the liver of catfish fed with different concentrations of proteins, lipids and carbohydrates. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 68(5): 1251-1258. <http://dx.doi.org/10.1590/1678-4162-8337>.
- MIRANDA, E.C.; GUIMARÃES, I.G.; CABRAL JUNIOR, C.R. 2009 Desempenho produtivo do tambaqui (*Colossoma macropomum*) alimentado com farinha de vagem de algaroba em substituição ao milho. *Pubvet*, 3: 1982-1263.
- MORI-PINEDO, L.A.; PEREIRA-FILHO, M.; OLIVEIRA-PEREIRA, M.I. 1999 Substituição do fubá de milho (*Zea mays* L.) por farinha de pupunha (*Bactris gasipaes*. H.B.K.) em ração para alevinos de tambaqui (*Colossoma macropomum*. Cuvier. 1818). *Acta Amazonica*, 23(3): 447-453. <http://dx.doi.org/10.1590/1809-43921999293453>.
- MOYANO, F.J.; CARDENETE, G.; DE LA HIGUERA, M. 1991 Nutritive and metabolic utilization of proteins with glutamic acid content by the rainbow trout *Oncorhynchus mykiss*. *Comparative Biochemistry and Physiology*, 100(3): 759-762. [http://dx.doi.org/10.1016/0300-9629\(91\)90404-Z](http://dx.doi.org/10.1016/0300-9629(91)90404-Z). PMID:1664307.
- MUIN, H.; FATAH, N.N.A.; NOR, M.H.M.; RAZAK, S.A. 2013 Rice bran replacement in *Clarias gariepinus* fingerlings diets with *Pleurotus florida* Stalk. *Sains Malaysiana*, 42(8): 1109-1114.
- NAGAE, M.Y.; HAYASHI, C.; SOARES, C.M.; FURUYA, W.M. 2002 Inclusão do milheto (*Pennisetum americanum*) em rações para alevinos de piauçu (*Leporinus macrocephalus*). *Revista Brasileira de Zootecnia*, 31(5): 1875-1880. <http://dx.doi.org/10.1590/S1516-35982002000800001>.
- NAQVI, S.U.F.; ABBAS, G.; AHMD, N.; AHMAD, I.; SHAHZAD, M.K.; QAMAR, S.H. 2015 Effect of replacement of maize gluten with raice bran (5:0 and 1:4) feed supplement on fish growth in a composite culture. *Universal Journal of Agriculture Research*, 3(4): 125-129.

- NYINA-WAMWIZA, L.; WATHELET, B.; RICHIR, J.; ROLLIN, X.; KESTEMONT, P. 2010 Partial or total replacement of fish meal by local agricultural by-products in diets of juvenile African catfish (*Clarias gariepinus*): growth performance, feed efficiency and digestibility. *Aquaculture Nutrition*, 16(3): 237-247. <http://dx.doi.org/10.1111/j.1365-2095.2009.00658.x>.
- PEREIRA JUNIOR, G.P.; PEREIRA, E.M.O.; PEREIRA FILHO, M.; BARBOSA, P.S.; SHIMODA, E.; BRANDÃO, L.V. 2013 Performance of juvenile tambaqui (*Colossoma macropomum*, Cuvier, 1818) fed diets containing crueira manioc flour (*Manihot esculenta*, Crantz) in replacement of corn (*Zea mays*). *Acta Amazonica*, 43(2): 217-226. <http://dx.doi.org/10.1590/S0044-59672013000200013>.
- RÊGO, M.M.T.; NEIVA, J.N.M.; RÊGO, A.C.; CÂNDIDO, M.J.D.; CARNEIRO, M.S.S.; LÔBO, R.N.B. 2010 Chemical and bromatological characteristics of elephant grass silages containing a mango by-product. *Revista Brasileira de Zootecnia*, 39(1): 81-87. <http://dx.doi.org/10.1590/S1516-35982010000100011>.
- RENUKDAS, N.; ENGLE, C.; LOCHMANN, R.; LI, M.H.; AVERY, J.; TUCKER, C.S.; BOSWORTH, B. 2014 Performance of alternative diets containing solvent-extracted distillers dried grains with soluble compared to traditional diets for pond-raised channel catfish, *Ictalurus punctatus*, and hybrid catfish, *Ictalurus punctatus* x *Ictalurus furcatus*. *Journal of the World Aquaculture Society*, 45(3): 290-300. <http://dx.doi.org/10.1111/jwas.12123>.
- RODRIGUES, A.P.O. 2016 Nutrição e alimentação do tambaqui (*Colossoma macropomum*). *Boletim do Instituto de Pesca*, 40(1): 135-145.
- SAMPAIO-OLIVEIRA, A.M.B.M.; CYRINO, J.E.P. 2008 Digestibility of plant protein-based diets by largemouth bass *Micropterus salmoides*. *Aquaculture Nutrition*, 14(4): 318-323. <http://dx.doi.org/10.1111/j.1365-2095.2007.00533.x>.
- SIGNOR, A.A.; BOSCOLO, W.R.; FEIDEN, A.; SIGNOR, A.; REIDEL, A. 2007 Triguilho na alimentação da tilápia do nilo (*Oreochromis niloticus* L.): digestibilidade e desempenho. *Ciência Rural*, 37(4): 1116-1121. <http://dx.doi.org/10.1590/S0103-84782007000400032>.
- SILVA, J.A.M.; PEREIRA-FILHO, M.; OLIVEIRA-PEREIRA, M.I. 2003 Frutos e sementes consumidos pelo tambaqui *Colossoma macropomum* (Cuvier, 1818) incorporados em rações. Digestibilidade e velocidade de trânsito pelo trato gastrointestinal. *Revista Brasileira de Zootecnia*, 32(6): 1815-1824. <http://dx.doi.org/10.1590/S1516-35982003000800003>.
- SILVA, P.C.; FRANÇA, A.F.; PADUA, D.M.C.; JACOB, G. 1997 *Milheto* (*Pennisetum americanum*) como substituto do milho (*Zea mays*) na alimentação do tambaqui (*Colossoma macropomum*). *Boletim do Instituto de Pesca*, 24(1): 125-131.
- SIPAÚBA-TAVARES, L.H.S. 1995 *Limnologia aplicada à aquicultura*. Jaboticabal: Funep. 72p.
- SKIBA-CASSY, S.; PANSEERAT, S.; LARQUIER, M.; DIAS, K.; SURGET, A.; PLAGNES-JUAN, E.; KAUSHIK, S.; SEILIEZ, I. 2013 Apparent low ability of liver and muscle to adapt to variation of dietary carbohydrate: protein ratio in rainbow trout (*Oncorhynchus mykiss*). *British Journal of Nutrition*, 109(8): 1359-1372. <http://dx.doi.org/10.1017/S0007114512003352>. PMID:22951215.
- SOUZA, R.C.; MELO, J.F.B.; NOGUEIRA FILHO, R.M.; CAMPECHE, D.F.B.; FIGUEIREDO, R.A.C.R. 2013 Influência da farinha de manga no crescimento e composição corporal da tilápia do Nilo. *Archivos de Zootecnia*, 62(238): 217-225. <http://dx.doi.org/10.4321/S0004-05922013000200007>.
- SOUZA, S.A.; SOUZA, R.C.; CAMPECHE, D.F.B.; CAMPOS, R.M.L.; MELO, J.F.B. 2014 Relação proteína: carboidrato no desempenho e no metabolismo de híbridos de *Pseudoplatystoma fasciatum* (fêmea) X *Leiarius marmoratus* (macho). *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 66(3): 879-886. <http://dx.doi.org/10.1590/1678-41625954>.
- ULLOAROJAS, J.B.; VERRETH, J.A.J. 2003 Growth of *Oreochromis aureus* fed with diets containing graded levels of coffee pulp and reared in two culture systems. *Aquaculture*, 217(1-4): 275-283. [http://dx.doi.org/10.1016/S0044-8486\(02\)00273-9](http://dx.doi.org/10.1016/S0044-8486(02)00273-9).
- XAVIER, B.; SAHU, N.P.; PAL, A.K.; JAIN, K.K.; MISRA, S.; DALVI, R.S.; BARUAH, K. 2012 Water soaking and exogenous enzyme treatment of plant based diets: effect on growth performance, whole-body composition and digestive enzymes activities of rohu, *Labeo rohita* (Hamilton), fingerlings. *Fish Physiology and Biochemistry*, 38(2): 341-353. <http://dx.doi.org/10.1007/s10695-011-9511-2>. PMID:21611720.